



Year: 2013

Dietary intake of vitamin D and calcium and breast cancer risk in the European Prospective Investigation into Cancer and Nutrition

Abbas, Sascha ; Linseisen, Jakob ; Rohrmann, Sabine ; Chang-Claude, Jenny ; Peeters, Petra H ; Engel, Pierre ; Brustad, Magritt ; Lund, Eiliv ; Skeie, Guri ; Olsen, Anja ; Tjønneland, Anne ; Overvad, Kim ; Boutron-Ruault, Marie-Christine ; Clavel-Chapelon, Francoise ; Fagherazzi, Guy ; Kaaks, Rudolf ; Boeing, Heiner ; Buijsse, Brian ; Adarakis, George ; Ouranos, Vassilis ; Trichopoulou, Antonia ; Masala, Giovanna ; Krogh, Vittorio ; Mattiello, Amalia ; Tumino, Rosario ; Sacerdote, Carlotta ; Buckland, Genevieve ; Suárez, Marcial Vicente Argüelles ; Sánchez, Maria-José ; Chirlaque, Maria-Dolores ; Barricarte, Aurelio ; Amiano, Pilar ; Manjer, Jonas ; Wirfält, Elisabet ; Lenner, Per ; Sund, Malin ; Bueno-de-Mesquita, H B ; van Duijnhoven, Fränzel J B ; Khaw, Kay-Tee ; Wareham, Nick ; Key, Timothy J ; Fedirko, Veronika ; Romieu, Isabelle ; Gallo, Valentina ; Norat, Teresa ; Wark, Petra A ; Riboli, Elio

Abstract: Studies assessing the effects of vitamin D or calcium intake on breast cancer risk have been inconclusive. Furthermore, few studies have evaluated them jointly. This study is the largest so far examining the association of dietary vitamin D and calcium intake with breast cancer risk in the European Prospective Investigation into Cancer and Nutrition. During a mean follow-up of 8.8 yr, 7760 incident invasive breast cancer cases were identified among 319,985 women. Multivariable Cox proportional hazards regression was used to estimate hazard ratios (HR) and 95% confidence intervals (CI) for pre- and postmenopausal breast cancer risk. Comparing the highest with the lowest quintile of vitamin D intake, HR and 95% CI were 1.07 (0.87-1.32) and 1.02 (0.90-1.16) for pre- and postmenopausal women, respectively. The corresponding HR and 95% CIs for calcium intake were 0.98 (0.80-1.19) and 0.90 (0.79-1.02), respectively. For calcium intake in postmenopausal women, the test for trend was borderline statistically significant ($P(\text{trend}) = 0.05$). There was no significant interaction between vitamin D and calcium intake and cancer risk ($P(\text{interaction}) = 0.57$ and 0.22 in pre- and postmenopausal women, respectively). In this large prospective cohort, we found no evidence for an association between dietary vitamin D or calcium intake and breast cancer risk.

DOI: <https://doi.org/10.1080/01635581.2013.752018>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-88647>

Journal Article

Accepted Version

Originally published at:

Abbas, Sascha; Linseisen, Jakob; Rohrmann, Sabine; Chang-Claude, Jenny; Peeters, Petra H; Engel, Pierre; Brustad, Magritt; Lund, Eiliv; Skeie, Guri; Olsen, Anja; Tjønneland, Anne; Overvad, Kim; Boutron-Ruault, Marie-Christine; Clavel-Chapelon, Francoise; Fagherazzi, Guy; Kaaks, Rudolf; Boeing, Heiner; Buijsse, Brian; Adarakis, George; Ouranos, Vassilis; Trichopoulou, Antonia; Masala, Giovanna;

Krogh, Vittorio; Mattiello, Amalia; Tumino, Rosario; Sacerdote, Carlotta; Buckland, Genevieve; Suárez, Marcial Vicente Argüelles; Sánchez, Maria-José; Chirlaque, Maria-Dolores; Barricarte, Aurelio; Amano, Pilar; Manjer, Jonas; Wirfält, Elisabet; Lenner, Per; Sund, Malin; Bueno-de-Mesquita, H B; van Duijnhoven, Fränzel J B; Khaw, Kay-Tee; Wareham, Nick; Key, Timothy J; Fedirko, Veronika; Romieu, Isabelle; Gallo, Valentina; Norat, Teresa; Wark, Petra A; Riboli, Elio (2013). Dietary intake of vitamin D and calcium and breast cancer risk in the European Prospective Investigation into Cancer and Nutrition. *Nutrition and Cancer*, 65(2):178-187.
DOI: <https://doi.org/10.1080/01635581.2013.752018>

Dietary intake of vitamin D and calcium and breast cancer risk in the European Prospective Investigation into Cancer and Nutrition

Sascha Abbas^{1,2}, Jakob Linseisen^{1,3}, Sabine Rohrmann^{1,4}, Jenny Chang-Claude¹, Petra H Peeters⁵, Pierre Engel⁶, Magritt Brustad⁷, Eiliv Lund⁷, Guri Skeie⁷, Anja Olsen⁸, Anne Tjønneland⁸, Kim Overvad⁹, Marie-Christine Boutron-Ruault⁶, Francoise Clavel-Chapelon⁶, Guy Fagherazzi⁶, Rudolf Kaaks¹, Heiner Boeing¹⁰, Brian Buijsse¹⁰, George Adarakis¹¹, Vassilis Ouranos¹¹, Antonia Trichopoulou^{11,12}, Giovanna Masala¹³, Vittorio Krogh¹⁴, Amalia Mattiello¹⁵, Rosario Tumino¹⁶, Carlotta Sacerdote^{17,18}, Genevieve Buckland¹⁹, Marcial Vicente Argüelles Suárez²⁰, Maria-José Sánchez^{21,22}, Maria-Dolores Chirlaque^{22,23}, Aurelio Barricarte^{22,24}, Pilar Amiano^{22,25}, Jonas Manjer²⁶, Elisabet Wirfält²⁷, Per Lenner²⁸, Malin Sund²⁹, H.B(as). Bueno-de-Mesquita^{30,31}, Fränzel J.B. van Duijnhoven^{30,32}, Kay-Tee Khaw³³, Nick Wareham³⁴, Timothy J Key³⁵, Veronika Fedirko³⁶, Isabelle Romieu³⁶, Valentina Gallo³⁷, Teresa Norat³⁷, Petra A Wark³⁷, Elio Riboli³⁷

¹ German Cancer Research Center, Division of Cancer Epidemiology, Heidelberg, Germany

² PMV Research Group at the Department of Child and Adolescent Psychiatry and Psychotherapy, University of Cologne, Cologne, Germany

³ Helmholtz Zentrum München, Institute of Epidemiology, Neuherberg, Germany

⁴ Institute of Social and Preventive Medicine, University of Zurich, Switzerland

⁵ Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands

⁶ Inserm Team 9 U1018 Center for Research in Epidemiology and Population Health, Team 9 and Paris South University, Institut Gustave-Roussy, Villejuif, France

⁷ Department of Community Medicine, University of Tromsø, Tromsø, Norway

⁸ Danish Cancer Society Research Center, Copenhagen, Denmark

⁹ Department of Epidemiology, School of Public Health, Aarhus University, Aarhus, Denmark

- ¹⁰ Department of Epidemiology, German Institute of Human Nutrition (DIfE) Potsdam-Rehbrücke, Nuthetal, Germany
- ¹¹ Hellenic Health Foundation, Athens, Greece
- ¹² WHO Collaborating Center for Food and Nutrition Policies, Department of Hygiene, Epidemiology and Medical Statistics, University of Athens Medical School, Athens, Greece
- ¹³ Molecular and Nutritional Epidemiology Unit, Cancer Research and Prevention Institute – ISPO, Florence, Italy
- ¹⁴ Nutritional Epidemiology Unit, Fondazione IRCCS Istituto Nazionale Tumori, Milan, Italy
- ¹⁵ Department of Clinical and Experimental Medicine, Federico II University, Naples, Italy
- ¹⁶ Cancer Registry and Histopathology Unit, ‘Civile-M.P. Arezzo’ Hospital, Ragusa, Italy
- ¹⁷ Human Genetics Foundation, Turin, Italy
- ¹⁸ CPO-Piemonte, Turin, Italy
- ¹⁹ Unit of Nutrition, Environment and Cancer, Cancer Epidemiology Research Programme, Catalan Institute of Oncology (ICO-IDIBELL), Barcelona, Spain
- ²⁰ Public Health Directorate, Asturias, Spain
- ²¹ Andalusian School of Public Health, Granada, Spain
- ²² CIBER Epidemiologia y Salud Publica (CIBERESP), Spain
- ²³ Department of Epidemiology, Murcia Regional Health Authority, Murcia, Spain
- ²⁴ Navarre Public Health Institute, Pamplona, Spain
- ²⁵ Public Health Division of Gipuzkoa, IIS Institute BioDonostia, Basque Health Department, San Sebastian, Spain
- ²⁶ Department of Plastic Surgery, Skane University Hospital, Malmö, Lund University, Malmö, Sweden
- ²⁷ Department of Clinical Sciences in Malmö, Malmö, Sweden
- ²⁸ Department of Radiation Sciences, Oncology, Umea University, Umea, Sweden
- ²⁹ Department of Surgical and Perioperative Sciences, Umea University, Umea, Sweden
- ³⁰ National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands,
- ³¹ Department of Gastroenterology and Hepatology, University Medical Center, Utrecht, The Netherlands.

³² Division of Human Nutrition, Wageningen University, Wageningen, The Netherlands

³³ Clinical Gerontology Unit, University of Cambridge, Cambridge, UK

³⁴ Medical Research Council Epidemiology Unit, Institute of Metabolic Science, University of Cambridge, Cambridge, UK

³⁵ Cancer Epidemiology Unit, Nuffield Department of Clinical Medicine, University of Oxford, Oxford, UK

³⁶ Nutritional Epidemiology Group, Nutrition and Metabolism Section, International Agency for Research on Cancer (IARC), Lyon, France

³⁷ Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK

Abstract

Studies assessing the effects of vitamin D or calcium intake on breast cancer risk have been inconclusive. Furthermore, few studies have evaluated them jointly. This study is the largest so far examining the association of dietary vitamin D and calcium intake with breast cancer risk in the European Prospective Investigation into Cancer and Nutrition.

During a mean follow-up of 8.8 years, 7,760 incident invasive breast cancer cases were identified among 319,985 women. Multivariable Cox proportional hazards regression was used to estimate hazard ratios (HR) and 95% confidence intervals (CI) for pre- and postmenopausal breast cancer risk.

Comparing the highest with the lowest quintile of vitamin D intake, HR and 95% CIs were 1.07 (0.87-1.32) and 1.02 (0.90-1.16) for pre- and postmenopausal women, respectively. The corresponding HR and 95% CIs for calcium intake were 0.98 (0.80-1.19) and 0.90 (0.79-1.02), respectively. For calcium intake in postmenopausal women, the test for trend was borderline statistically significant ($p_{trend}=0.05$). There was no significant interaction between vitamin D and calcium intake and cancer risk ($p_{interaction}=0.57$ and 0.22 in pre- and postmenopausal women, respectively).

In this large prospective cohort, we found no evidence for an association between dietary vitamin D or calcium intake and breast cancer risk.

Keywords

Vitamin D intake, calcium intake, cohort study, breast cancer risk, diet

Introduction

Both epidemiological and experimental data suggest an inverse association between vitamin D and breast cancer (1-3). Besides its classical function on calcium homeostasis and bone metabolism, vitamin D has potential anticarcinogenic properties with regard to regulation of apoptosis, cell differentiation, cell growth and growth factor signaling (4-6).

Studies assessing the association of vitamin D intake and breast cancer risk have yielded inconsistent results. A meta-analysis from 2010 that pooled 11 studies assessing vitamin D intake from either diet and/or supplements and breast cancer risk resulted in a significant overall association with a RR of 0.91 (0.85 – 0.97) (7). However, there was no significant association when assessing vitamin D intake from diet only. Another meta-analysis from 2008 reported no statistically significant association between dietary vitamin D intake from food or supplements and breast cancer risk (8). A number of studies have assessed the relationship between breast cancer risk and serum vitamin D metabolites [25(OH)D or 1,25(OH)₂D], the biomarkers for vitamin D status in humans, which cover vitamin D from diet, supplements and from endogenous production after exposure to sunlight. Most case-control studies reported an inverse association between vitamin D status and breast cancer risk (9). However, there was no statistically significant inverse association in cohort studies (9), except for a recent study in the French E3N Cohort (10). In addition, the Second Expert Report by the World Cancer Research Fund (WCRF) classified the evidence as limited (no conclusion) for both pre- and postmenopausal breast cancer (11).

There is less scientific evidence with regard to the anticarcinogenic potential of calcium, though effects of calcium on cell proliferation and apoptosis have been reported (12, 13). However, due to the tight homeostasis of plasma calcium, it is unclear whether dietary calcium may affect breast cancer risk. Various studies assessed the association between dietary calcium intake and breast cancer risk with the majority reporting inverse associations (3). A meta-analysis pooling data from 6 cohort and 9 case-control studies resulted in a

statistically significant inverse association between calcium intake and breast cancer risk. However, the results suggest an inverse association in pre- but not in postmenopausal women (7).

Vitamin D intake varies considerably across countries, which may also lead to different risk estimates (8). Furthermore, the amount of dietary calcium intake may affect the association between dietary vitamin D intake and breast cancer risk. We report here on the association between dietary vitamin D and calcium intake from foods and breast cancer risk in a heterogeneous population from different countries in Europe within the European Prospective Investigation into Cancer and Nutrition (EPIC). Due to its large sample size, EPIC provides the opportunity to investigate the association for both pre- and postmenopausal breast cancer risk. In addition to main effects, potential interaction effects between dietary vitamin D and calcium intake are addressed.

Material and methods

Study population

EPIC is a large prospective cohort study conducted since 1992 in 23 centers in 10 European countries [Denmark (Aarhus, Copenhagen), France, Germany (Heidelberg, Potsdam), Great Britain (Cambridge, Oxford), Greece, Italy (Florence, Varese, Ragusa, Turin, Naples), Norway, Spain (Asturias, Granada, Murcia, Navarra, San Sebastian), Sweden (Malmo, Umea), The Netherlands (Bilthoven, Utrecht)] with over 500,000 participants (14). Generally, the centers recruited the participants from the general population. However, French participants were members of a health insurance for school and university employees. Spain and Italy recruited participants among blood donors, members of several health insurance programs, employees of several enterprises, civil servants but also the general population. Participants were recruited from members of screening programs in Utrecht and Florence. In France, Norway, Utrecht, and Naples only women were included. Participants from Greece

were excluded due to missing validated information on vitamin D and calcium intake. Of the 342,686 female participants, we excluded 6,822 subjects with missing follow-up information, 15,019 subjects without dietary information, 837 subjects with in situ breast cancer, and 23 with not clearly malignant breast tumors (e.g. phyllodes tumors). Thus, the final cohort consisted of 319,985 participants.

Exposure assessment

Diet over the previous twelve months was assessed using dietary assessment instruments that were specifically developed for each participating country (14). The questions were structured by common food groups except for the questionnaires used in Italy and Spain, where questions were structured by country-specific meals.

All participants were asked to report their average consumption of each food item over the previous twelve months by structured categories ranging from never or less than once per month to six or more times per day. All dietary measurement instruments have been validated previously in a series of studies within the various source populations participating in EPIC (15).

To collect information on education, medical history (surgeries and previous illnesses), tobacco and alcohol consumption, and physical activity and other lifestyle factors, a further questionnaire was used. Height and weight were measured at baseline, except for most participants recruited at the Oxford study center, and in France as well as all participants recruited in Norway. Self-reported data were available for the latter centres (14).

Menopausal status at baseline was defined as described previously using an algorithm that accounts for information on menstrual status/history, type of menopause, use of oral contraceptives and menopausal hormones (16).

Outcome assessment

In most countries (Denmark, Italy, The Netherlands, Norway, Spain, Sweden, and the UK) incident cancers were identified through linkage with population registries. Cancer cases were further identified by active follow-up, through a health insurance company (France) or direct contacts with study subjects, their doctors or their next of kin (France, Germany, and Greece). Mortality data were also obtained from either the cancer or mortality registries at the regional or national level. Participants from centers that relied on registry data that have been reported to IARC were censored between December 2002 and December 2005, depending on the study center. For Germany and Greece, the end of the follow up was considered to be the last known contact, the date of diagnosis, or the date of death, whichever came first. Cancer incidence was coded according to ICD-O-2.

Statistical analysis

Participants were categorized according to quintiles of dietary intake of vitamin D and calcium. Cox proportional hazards regression was used to examine the association of dietary intake of vitamin D or calcium and breast cancer risk. Age was used as the primary time variable in the Cox models. Time at entry was age at recruitment, exit time was age when participants were diagnosed with cancer, died, were lost to follow-up, or were censored at the end of the follow-up period, whichever came first. Analyses were stratified by center to account for center effects such as follow-up procedures and questionnaire design, and by 1-year categories of age at recruitment to account for possible violations of the proportional hazard assumption.

Different models were run to fully disentangle the effects of dietary vitamin D and calcium on breast cancer risk: model 1, energy-adjusted only (kcal/day, quintiles); model 2, further adjustment for potential confounders and risk factors defined a-priori: total energy excluding energy from fat and alcohol (kcal/day, quintiles), fat (g/day, quintiles, monounsaturated,

polyunsaturated and saturated fat), alcohol consumption (g/day, quintiles), weight (kg, cont.), height (cm, cont.), smoking status (never, former, current, unknown), education level (none or primary school, technical/professional school, secondary school, longer education incl. university degree, unknown), menopausal status (premenopausal, perimenopausal, postmenopausal), total physical activity (inactive, moderately inactive, moderately active, active, missing), current use of oral contraceptives or hormones for menopause (yes, no, unknown), age at menarche (d12, 13-14, e15 years, missing); model 3, further adjustment on vitamin D ($\mu\text{g/day}$, quintiles) and calcium intake (mg/day, quintiles) in the calcium and vitamin D models, respectively. We adjusted for non-fat, non-alcohol energy and independently adjusted for alcohol and fat to account for both nutrients as potential confounders. In a sensitivity analysis we recalculated the models for all, pre- and postmenopausal women using the complete case analysis.

Multiplicative statistical interaction was evaluated with the likelihood-ratio-test by including cross-product terms of the potential interaction variables (continuous intake variables for vitamin D and calcium) in the respective models. Tests for trend were performed using the Wald statistic of the continuous intake variable.

In order to compare results across populations with different levels of sunlight exposure, the EPIC centres were divided into 7 latitude groups: below 42°N (Granada, Murcia, Ragusa, and Naples); 42°N - 44°N (Asturias, Navarra, San Sebastian, Florence, and South coast of France – centred in Marseille); 45°N - 46°N (Varese, Turin, and South of France – centred in Lyon); 47°N - 49°N (North-East and West of France – centred in Nantes and Paris, respectively – and Heidelberg); 50°N - 51°N (Potsdam, Utrecht, Bilthoven, Cambridge, and Oxford); 52°N - 56°N (Malmö, Aarhus, and Copenhagen); and above 57°N (Norway, and Umeå)

To correct for measurement error, FFQ dietary intake data were calibrated against 24-hour dietary recall data obtained from a random sample of the subcohort (N = 36,994). A fixed-effects linear model was used in which center and sex-specific 24-h recall data were regressed

on the FFQ intakes (17-19). The calibrated FFQ data were used to model dietary intake of vitamin D or calcium continuously. All tests were two-sided and considered to be statistically significant with a p-value of < 0.05 . All analyses were performed in SAS software version 9.2 (Cary, NC, US).

Results

Mean follow-up time for the study population was 8.8 years. Mean time from baseline until breast cancer diagnosis was 5.1 years. During the follow-up, a total of 7,760 incident invasive (primary, malignant) breast cancer cases were identified, including 1,802 women who were premenopausal at recruitment and 4,259 cases in women who were postmenopausal at recruitment. Descriptive characteristics (at baseline) of the total cohort of 319,985 women are shown in Table 1. Women in the highest quintile of both vitamin D and calcium intake had a higher non-fat energy intake and a higher fat intake as compared to women in the lowest quintile. Furthermore, women in the lower quintile of vitamin D intake weighed less, used oral contraceptives less frequently, were younger at menarche, and were more often non-smokers.

As shown in Table 2, dietary vitamin D intake was highest in EPIC Sweden as compared to the other countries; calcium intake was highest in Denmark and the Netherlands.

Dietary vitamin D intake was not associated with overall breast cancer risk ($p_{trend} = 0.92$; Table 3), There was also no association when addressing pre- or postmenopausal women separately.

We found some indication for an inverse association between dietary calcium intake and breast cancer risk. The hazard ratio (HR) (95% confidence interval (CI)) for the highest as compared to the lowest quintile was 0.91 (0.83–1.01) (Table 3). The test for trend did not

reach statistical significance ($p_{trend} = 0.06$). Stratification by menopausal status indicated a possible association in postmenopausal women 0.90 (0.79–1.02), ($p_{trend} = 0.05$).

In the complete case analysis results were similar to those when using confounders with a category for missing data for both dietary vitamin D and calcium intake.

As dietary vitamin D may modify the association between calcium intake and breast cancer risk (and vice versa), we assessed potential interaction effects between both nutrients. However, statistical significant interaction ($p_{interaction} = 0.44$; 0.57 and 0.22 for all, premenopausal and postmenopausal women, respectively; data not shown) was not observed. Moreover, additional adjustment by vitamin D ($\mu\text{g/day}$, quintiles) and calcium intakes (mg/day , quintiles) in the calcium and vitamin D models, respectively, did not result in changes of the risk estimates (data not shown).

To correct for potential measurement error, we repeated the analysis with the questionnaire data calibrated against 24-h recall data for the continuous intake variables. The overall associations were 1.00 (95% CI: 0.99-1.01; $p = 0.92$; uncalibrated) and 1.02 (95% CI: 0.99-1.06; $p = 0.22$; calibrated) per 1 $\mu\text{g/day}$ increment of dietary vitamin D and 0.99 (95% CI: 0.99-1.00; $p = 0.06$; uncalibrated) and 0.99 (95% CI: 0.97-1.00; $p = 0.11$; calibrated) per 100 mg/day increment of calcium intake.

In a country-specific analysis with the continuous calibrated intake values there was no statistically significant association between either vitamin D or calcium intake and overall breast cancer risk. Country-specific results by menopausal status also showed no significant associations except for an increased breast cancer risk associated with calcium intake in premenopausal women in the Netherlands ($p = 0.01$) (data not shown).

As a proxy for sun exposure and thus endogenous vitamin D production we divided the centres in 7 categories according to their latitude. Adjustment for latitude did not change the observed risk estimates for vitamin D and breast cancer risk. Moreover, we calculated

separate risk estimates for each latitude. There was no significant association between vitamin D and breast cancer risk in any of the latitude groups ($p_{\text{heterogeneity}} = 0.37$).

Discussion

The present study is so far the largest prospective study including data from nine European countries that assessed dietary vitamin D or calcium intake and breast cancer risk. Dietary vitamin D and calcium intake was not associated with breast cancer risk. There was also no significant interaction between both nutrients. Evidence for an association between dietary vitamin D intake and breast cancer risk has not been consistent in cohort studies, which reported a significant inverse association only in premenopausal women (20), in postmenopausal women with ER positive tumors (21) or no significant association at all (22-28). Four out of eight case-control studies, three of which included premenopausal women only, did not find an association (29-33), three reported significantly inverse associations (34-36), whereas one study reported a non-significantly increased risk with higher dietary vitamin D intake (37). A meta-analysis that pooled results from 11 studies assessing total vitamin D intake and breast cancer risk resulted in a significant overall inverse association with risk of 0.91 (0.85 – 0.97) (7). However, when differentiated by source of vitamin D, the significant inverse association was seen in vitamin D supplement users only but no association existed when analysing dietary vitamin D intake from foods. Also, a meta-analysis from 2008 reported no significant association between dietary vitamin D intake from food or supplements and breast cancer risk (8). However, the authors observed a significant inverse association when restricting the analysis to women taking more than 10µg/day vitamin D. Thus, the amount of dietary intake in the present population may be too low to detect a potential effect. Intake of dietary vitamin D in the present study was somehow comparable with dietary intake, but not total intake values reported in US studies (20, 21). In contrast,

dietary calcium intake in EPIC is somehow higher than reported dietary intake in the US (20, 21). Unfortunately, we did not have information on supplemental vitamin D intake nor on sun exposure as primary source of vitamin D. We, however, adjusted in the regression analysis for physical activity which may in part account for sun exposure, with a marginal effect on risk estimates. In a sensitivity analysis, we also adjusted for centers' latitude as another proxy for sun exposure, which also did not affect the risk estimates. Furthermore analyses were stratified by centres latitude which may in part account for differential effects of sun exposure by region. In contrast to Engel et al. (26), who observed a decreased postmenopausal breast cancer risk associated with vitamin D intake in regions with high ultraviolet radiation dose, we did not find any evidence for a differential effect of dietary vitamin D on pre- or postmenopausal breast cancer risk in regions with different sun exposure.

Studies accounting for total vitamin D status by measuring 25(OH)D in plasma or serum have yielded mostly significant inverse associations in case-control studies but, except for one study (10), no associations with risk in cohort studies (9).

We did not observe a statistically significant association between dietary calcium and breast cancer risk in pre- or postmenopausal women. However, we found some indication of an inverse association in postmenopausal women with borderline significance ($p_{trend} = 0.05$).

Most case-control studies reported an inverse association (37-46), although statistically significant only in a subset (40, 41, 45, 46) and no association was observed in three studies in premenopausal women only (30, 31, 34). Cohort studies reported a reduced risk of breast cancer associated with dietary calcium (20, 21, 24, 47, 48), including two studies in premenopausal women only (20, 24) and one in estrogen receptor-/progesterone receptor-negative tumors only (48), or no significant association (33, 49). The results of the largest trial so far, the Women's Health Initiative, did not demonstrate an inverse association between breast cancer risk and calcium plus vitamin D supplementation (50). However, a recent reanalysis of the WHI data provided evidence for a protective effect of vitamin D and calcium

supplementation in women who were not taking further personal supplements, indicating that supplementation may only be effective in those women who were deficient (51).

Calcium might exert its potential anticarcinogenic properties via several mechanisms including effects on cell proliferation, apoptosis or cell differentiation (3, 12, 13). So far, differential anticarcinogenic mechanisms on pre- and postmenopausal breast cancer are not known. However, a meta-analysis pooling data from 15 studies reported a statistically significant inverse association in pre-/perimenopausal women but not postmenopausal women (7).

Further studies have concentrated on measurement of calcium in serum samples as a marker of a subject's calcium status (52-54). Results have been inconsistent, showing significant positive (53) or negative associations (52) in premenopausal women or no significant association at all (54). However, as the calcium metabolism and thus serum calcium levels are tightly regulated, blood levels of calcium may not be a good marker for calcium status or dietary calcium intake (55).

In line with previous studies, our results do not suggest an interaction between vitamin D and calcium intake (20, 33, 34), nor that the associations were confounded by each other. In contrast, another study reported a borderline significant interaction between both nutrients in postmenopausal women ($p = 0.05$) with a non-significant inverse association for calcium intake in the group with the highest tertile vitamin D intake (24).

Strengths of the study are its prospective design, the calibration of food frequency questionnaire data against 24-h-recall data to reduce measurement error, as well as the inclusion of participants from different European countries reflecting a wide range of intake.

However, several limitations have to be taken into account. We did not have information on the overall vitamin D status, including vitamin D from endogenous synthesis and vitamin D from supplements. Vitamin D supplementation varies across European countries. Especially

in the Northern countries, vitamin D from supplements, e.g. cod liver oil, is frequently used and thus one of the most important sources of vitamin D (56). Since the contribution of diet to the overall vitamin D status of a person is limited (not more than 10-20% of vitamin D may originate from dietary sources), studies with information on the entire vitamin D status may come to another conclusion. Moreover, we assessed nutrient intake at only one point in life time. It is currently unknown at what point in time vitamin D or calcium intake may affect breast cancer risk. However, a recent study measuring ultraviolet sunlight exposure as a proxy for vitamin D production suggest that exposure in both adolescence and adulthood may reduce breast cancer risk (57)

In summary, the present study does not support an association of dietary vitamin D or calcium intake with breast cancer risk. Further research assessing vitamin D and breast cancer risk should focus on biomarker measurement to account for the overall vitamin D status as dietary intake may be too low to detect potential anticarcinogenic effects

Institution where work was performed

German Cancer Research Center, Division of Cancer Epidemiology, Heidelberg, Germany

The manuscript has not been published and has not been submitted simultaneously for publication elsewhere.

Financial support

EPIC is financially supported by the European Commission (DG-SANCO) and the International Agency for Research on Cancer. The national cohorts are supported by Danish Cancer Society (Denmark); Ligue Contre le Cancer, Institut Gustave Roussy, Mutuelle

Générale de l'Education Nationale, Institut National de la Santé et de la Recherche Médicale (INSERM) (France); Deutsche Krebshilfe, Deutsches Krebsforschungszentrum and Federal Ministry of Education and Research (Germany); Hellenic Health Foundation, Stavros Niarchos Foundation and Hellenic Ministry of Health and Social Solidarity (Greece); Italian Association for Research on Cancer (AIRC) and National Research Council (Italy); Dutch Ministry of Public Health, Welfare and Sports (VWS), Netherlands Cancer Registry (NKR), LK Research Funds, Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research Fund (WCRF), Statistics Netherlands (The Netherlands); ERC-2009-AdG 232997 and Nordforsk, Nordic Center of Excellence programme on Food, Nutrition and Health. (Norway); Health Research Fund (FIS), Regional Governments of Andalucía, Asturias, Basque Country, Murcia (no. 6236) and Navarra, ISCIII RETIC (RD06/0020) (Spain); Swedish Cancer Society, Swedish Scientific Council and Regional Government of Skåne and Västerbotten (Sweden); Cancer Research UK, Medical Research Council (United Kingdom); The Catalan Institute of Oncology and RTICC 'Red Tematica de Investigacion Cooperativa en Cancer (R06/0020) (Spain).

References

Reference List

1. Colston KW and Hansen CM: Mechanisms implicated in the growth regulatory effects of vitamin D in breast cancer. *Endocrine-Related Cancer* **9**, 45-59, 2002.
2. Welsh J: Vitamin D and prevention of breast cancer. *Acta Pharmacol Sin* **28**, 1373-1382, 2007.
3. Cui Y and Rohan TE: Vitamin D, calcium, and breast cancer risk: a review. *Cancer Epidemiol Biomarkers Prev* **15**, 1427-1437, 2006.
4. Trump DL, Hershberger PA, Bernardi RJ, Ahmed S, Muindi J, et al.: Anti-tumor activity of calcitriol: pre-clinical and clinical studies. *J Steroid Biochem Mol Biol* **89-90**, 519-526, 2004.
5. Jensen SS, Madsen MW, Lukas J, Binderup L, and Bartek J: Inhibitory effects of 1alpha,25-dihydroxyvitamin D(3) on the G(1)-S phase-controlling machinery. *Mol Endocrinol* **15**, 1370-1380, 2001.

6. Dusso AS, Brown AJ, and Slatopolsky E: Vitamin D. *Am J Physiol Renal Physiol* **289**, F8-28, 2005.
7. Chen P, Hu P, Xie D, Qin Y, Wang F, et al.: Meta-analysis of vitamin D, calcium and the prevention of breast cancer. *Breast Cancer Res Treat* **121**, 469-477, 2010.
8. Gissel T, Rejnmark L, Mosekilde L, and Vestergaard P: Intake of vitamin D and risk of breast cancer--a meta-analysis. *J Steroid Biochem Mol Biol* **111**, 195-199, 2008.
9. Yin L, Grandi N, Raum E, Haug U, Arndt V, et al.: Meta-analysis: serum vitamin D and breast cancer risk. *Eur J Cancer* **46**, 2196-2205, 2010.
10. Engel P, Fagherazzi G, Boutten A, Dupre T, Mesrine S, et al.: Serum 25(OH) vitamin D and risk of breast cancer: a nested case-control study from the French E3N cohort. *Cancer Epidemiol Biomarkers Prev* **19**, 2341-2350, 2010.
11. World Cancer Research Fund / American Institute for Cancer Research: Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. Washington, DC: 2007.
12. Mathiasen IS, Sergeev IN, Bastholm L, Elling F, Norman AW, et al.: Calcium and calpain as key mediators of apoptosis-like death induced by vitamin D compounds in breast cancer cells. *J Biol Chem* **277**, 30738-30745, 2002.
13. Whitfield JF, Boynton AL, MacManus JP, Sikorska M, and Tsang BK: The regulation of cell proliferation by calcium and cyclic AMP. *Mol Cell Biochem* **27**, 155-179, 1979.
14. Riboli E, Hunt KJ, Slimani N, Ferrari P, Norat T, et al.: European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr* **5**, 1113-1124, 2002.
15. Margetts BM and Pietinen P: European Prospective Investigation into Cancer and Nutrition: validity studies on dietary assessment methods. *Int J Epidemiol* **26 Suppl 1**, S1-S5, 1997.
16. Lahmann PH, Hoffmann K, Allen N, van Gils CH, Khaw KT, et al.: Body size and breast cancer risk: findings from the European Prospective Investigation into Cancer And Nutrition (EPIC). *Int J Cancer* **111**, 762-771, 2004.
17. Slimani N, Kaaks R, Ferrari P, Casagrande C, Clavel-Chapelon F, et al.: European Prospective Investigation into Cancer and Nutrition (EPIC) calibration study: rationale, design and population characteristics. *Public Health Nutr* **5**, 1125-1145, 2002.
18. Slimani N, Ferrari P, Ocke M, Welch A, Boeing H, et al.: Standardization of the 24-hour diet recall calibration method used in the european prospective investigation into cancer and nutrition (EPIC): general concepts and preliminary results. *Eur J Clin Nutr* **54**, 900-917, 2000.
19. Ferrari P, Day NE, Boshuizen HC, Roddam A, Hoffmann K, et al.: The evaluation of the diet/disease relation in the EPIC study: considerations for the calibration and the disease models. *Int J Epidemiol* **37**, 368-378, 2008.

20. Shin MH, Holmes MD, Hankinson SE, Wu K, Colditz GA, et al.: Intake of dairy products, calcium, and vitamin d and risk of breast cancer. *J Natl Cancer Inst* **94**, 1301-1311, 2002.
21. McCullough ML, Rodriguez C, Diver WR, Feigelson HS, Stevens VL, et al.: Dairy, calcium, and vitamin D intake and postmenopausal breast cancer risk in the Cancer Prevention Study II Nutrition Cohort. *Cancer Epidemiol Biomarkers Prev* **14**, 2898-2904, 2005.
22. John EM, Schwartz GG, Dreon DM, and Koo J: Vitamin D and breast cancer risk: the NHANES I Epidemiologic follow-up study, 1971-1975 to 1992. National Health and Nutrition Examination Survey. *Cancer Epidemiol Biomarkers Prev* **8**, 399-406, 1999.
23. Frazier AL, Li L, Cho E, Willett WC, and Colditz GA: Adolescent diet and risk of breast cancer. *Cancer Causes Control* **15**, 73-82, 2004.
24. Lin J, Manson JE, Lee IM, Cook NR, Buring JE, et al.: Intakes of calcium and vitamin D and breast cancer risk in women. *Arch Intern Med* **167**, 1050-1059, 2007.
25. Kuper H, Yang L, Sandin S, Lof M, Adami HO, et al.: Prospective study of solar exposure, dietary vitamin D intake, and risk of breast cancer among middle-aged women. *Cancer Epidemiol Biomarkers Prev* **18**, 2558-2561, 2009.
26. Engel P, Fagherazzi G, Mesrine S, Boutron-Ruault MC, and Clavel-Chapelon F: Joint effects of dietary vitamin D and sun exposure on breast cancer risk: results from the French E3N cohort. *Cancer Epidemiol Biomarkers Prev* **20**, 187-198, 2011.
27. Edvardsen K, Veierod MB, Brustad M, Braaten T, Engelsen O, et al.: Vitamin D-effective solar UV radiation, dietary vitamin D and breast cancer risk. *Int J Cancer* **128**, 1425-1433, 2011.
28. Robien K, Cutler GJ, and Lazovich D: Vitamin D intake and breast cancer risk in postmenopausal women: the Iowa Women's Health Study. *Cancer Causes Control* **18**, 775-782, 2007.
29. Simard A, Vobecky J, and Vobecky JS: Vitamin D deficiency and cancer of the breast: an unprovocative ecological hypothesis. *Can J Public Health* **82**, 300-303, 1991.
30. Witte JS, Ursin G, Siemiatycki J, Thompson WD, Paganini-Hill A, et al.: Diet and premenopausal bilateral breast cancer: a case-control study. *Breast Cancer Res Treat* **42**, 243-251, 1997.
31. Potischman N, Swanson CA, Coates RJ, Gammon MD, Brogan DR, et al.: Intake of food groups and associated micronutrients in relation to risk of early-stage breast cancer. *Int J Cancer* **82**, 315-321, 1999.
32. Frazier AL, Ryan CT, Rockett H, Willett WC, and Colditz GA: Adolescent diet and risk of breast cancer. *Breast Cancer Res* **5**, R59-R64, 2003.
33. Anderson LN, Cotterchio M, Vieth R, and Knight JA: Vitamin D and calcium intakes and breast cancer risk in pre- and postmenopausal women. *Am J Clin Nutr* **91**, 1699-1707, 2010.

34. Abbas S, Linseisen J, and Chang-Claude J: Dietary vitamin d and calcium intake and premenopausal breast cancer risk in a german case-control study. *Nutr Cancer* **59**, 54-61, 2007.
35. Rossi M, McLaughlin JK, Laggiou P, Bosetti C, Talamini R, et al.: Vitamin D intake and breast cancer risk: a case-control study in Italy. *Ann Oncol* **20**, 374-378, 2009.
36. Blackmore KM, Lesosky M, Barnett H, Raboud JM, Vieth R, et al.: Vitamin D from dietary intake and sunlight exposure and the risk of hormone-receptor-defined breast cancer. *Am J Epidemiol* **168**, 915-924, 2008.
37. Levi F, Pasche C, Lucchini F, and La Vecchia C: Dietary intake of selected micronutrients and breast-cancer risk. *Int J Cancer* **91**, 260-263, 2001.
38. Braga C, La Vecchia C, Negri E, Franceschi S, and Parpinel M: Intake of selected foods and nutrients and breast cancer risk: an age- and menopause-specific analysis. *Nutr Cancer* **28**, 258-263, 1997.
39. Negri E, La Vecchia C, Franceschi S, D'Avanzo B, Talamini R, et al.: Intake of selected micronutrients and the risk of breast cancer. *Int J Cancer* **65**, 140-144, 1996.
40. Landa MC, Frago N, and Tres A: Diet and the risk of breast cancer in Spain. *Eur J Cancer Prev* **3**, 313-320, 1994.
41. Zaridze D, Lifanova Y, Maximovitch D, Day NE, and Duffy SW: Diet, alcohol consumption and reproductive factors in a case-control study of breast cancer in Moscow. *Int J Cancer* **48**, 493-501, 1991.
42. Boyapati SM, Shu XO, Jin F, Dai Q, Ruan Z, et al.: Dietary calcium intake and breast cancer risk among Chinese women in Shanghai. *Nutr Cancer* **46**, 38-43, 2003.
43. Adzersen KH, Jess P, Freivogel KW, Gerhard I, and Bastert G: Raw and cooked vegetables, fruits, selected micronutrients, and breast cancer risk: a case-control study in Germany. *Nutr Cancer* **46**, 131-137, 2003.
44. Katsouyanni K, Willett W, Trichopoulos D, Boyle P, Trichopoulou A, et al.: Risk of breast cancer among Greek women in relation to nutrient intake. *Cancer* **61**, 181-185, 1988.
45. Kawase T, Matsuo K, Suzuki T, Hirose K, Hosono S, et al.: Association between vitamin D and calcium intake and breast cancer risk according to menopausal status and receptor status in Japan. *Cancer Sci* **101**, 1234-1240, 2010.
46. Zhang CX, Ho SC, Fu JH, Cheng SZ, Chen YM, et al.: Dairy products, calcium intake, and breast cancer risk: a case-control study in china. *Nutr Cancer* **63**, 12-20, 2011.
47. Knekt P, Jarvinen R, Seppanen R, Pukkala E, and Aromaa A: Intake of dairy products and the risk of breast cancer. *Br J Cancer* **73**, 687-691, 1996.
48. Larsson SC, Bergkvist L, and Wolk A: Long-term dietary calcium intake and breast cancer risk in a prospective cohort of women. *Am J Clin Nutr* **89**, 277-282, 2009.

49. Hjartaker A, Thoresen M, Engeset D, and Lund E: Dairy consumption and calcium intake and risk of breast cancer in a prospective cohort: The Norwegian Women and Cancer study. *Cancer Causes Control*, 2010.
50. Chlebowski RT, Johnson KC, Kooperberg C, Pettinger M, Wactawski-Wende J, et al.: Calcium plus vitamin D supplementation and the risk of breast cancer. *J Natl Cancer Inst* **100**, 1581-1591, 2008.
51. Bolland MJ, Grey A, Gamble GD, and Reid IR: Calcium and vitamin D supplements and health outcomes: a reanalysis of the Women's Health Initiative (WHI) limited-access data set. *Am J Clin Nutr* **94**, 1144-1149, 2011.
52. Almquist M, Manjer J, Bondeson L, and Bondeson AG: Serum calcium and breast cancer risk: results from a prospective cohort study of 7,847 women. *Cancer Causes Control* **18**, 595-602, 2007.
53. Almquist M, Bondeson AG, Bondeson L, Malm J, and Manjer J: Serum levels of vitamin D, PTH and calcium and breast cancer risk-a prospective nested case-control study. *Int J Cancer* **127**, 2159-2168, 2010.
54. Cui Y, Vogt S, Olson N, Glass AG, and Rohan TE: Levels of zinc, selenium, calcium, and iron in benign breast tissue and risk of subsequent breast cancer. *Cancer Epidemiol Biomarkers Prev* **16**, 1682-1685, 2007.
55. Nordin BC: Calcium requirement is a sliding scale. *Am J Clin Nutr* **71**, 1381-1383, 2000.
56. Skeie G, Braaten T, Hjartaker A, Lentjes M, Amiano P, et al.: Use of dietary supplements in the European Prospective Investigation into Cancer and Nutrition calibration study. *Eur J Clin Nutr* **63 Suppl 4**, S226-S238, 2009.
57. Anderson LN, Cotterchio M, Kirsh VA, and Knight JA: Ultraviolet sunlight exposure during adolescence and adulthood and breast cancer risk: a population-based case-control study among Ontario women. *Am J Epidemiol* **174**, 293-304, 2011.

Table 1: Descriptive characteristics of the EPIC cohort according to the lowest and highest vitamin D and calcium intake

	Total cohort	Dietary vitamin D intake ³		Dietary calcium intake ³	
		Lowest quintile	Highest quintile	Lowest quintile	Highest quintile
Age (years), mean (SD)	50.2 (9.7)	48.9 (10.7)	50.3 (9.4)	49.1 (8.9)	50.9 (9.9)
Height (cm), mean (SD)	162.6 (6.6)	161.0 (6.6)	164.7 (6.3)	162.5 (6.7)	163.1 (6.5)
Weight (kg), mean (SD)	65.5 (11.6)	63.1 (11.2)	67.7 (11.8)	65.8 (11.7)	65.9 (11.8)
Ethanol intake (g/d), mean (SD)	8.3 (11.8)	7.7 (11.8)	6.7 (10.1)	6.9 (11.1)	9.1 (12.3)
Non-fat energy intake (kcal/d), mean (SD)	1196 (342)	1079 (317)	1263 (330)	919 (233)	1496 (340)
FAMS (g/d), mean (SD)	27.4 (11.1)	23.8 (10.1)	28.7 (10.5)	20.8 (8.1)	34.3 (12.6)
FAPU (g/d), mean (SD)	13.5 (5.9)	11.8 (5.9)	15.6 (5.9)	10.9 (4.8)	16.0 (6.6)
FASAT (g/d), mean (SD)	29.7 (11.8)	23.9 (9.7)	33.0 (12.4)	20.9 (7.5)	39.8 (13.1)
Education ¹					
Primary school completed (%)	24.0	25.3	25.4	29.0	20.6
University degree (%)	23.0	27.4	17.6	18.7	25.1
Missing (%)	4.5	5.5	3.0	3.3	4.7
Smoking status					
Never (%)	54.8	62.3	44.9	49.2	56.2
Former (%)	23.4	20.4	26.5	23.2	24.2
Current (%)	19.6	15.4	26.0	25.1	17.5
Missing (%)	2.2	1.9	2.5	2.5	2.1
Age at menarche					
d12 years (%)	35.2	39.4	29.4	33.5	35.7
13-14 years (%)	46.1	46.2	47.6	46.6	45.8
e15 years (%)	15.2	13.3	18.0	15.7	15.7
Missing (%)	3.5	1.2	5.0	4.2	2.9
Current use of oral contraceptives/hormones for menopause					
Yes	22.1	20.0	23.7	22.9	21.8
No	71.4	76.1	68.2	69.8	72.7
Missing (%)	6.5	3.9	8.1	7.4	5.6
Physical activity ²					
Inactive (%)	12.2	14.5	7.2	10.7	12.0
Moderately inactive (%)	29.6	36.4	17.4	22.0	33.3

Moderately active (%)	34.6	39.4	26.9	28.4	38.7
Active (%)	7.6	7.1	7.2	5.1	10.9
Missing (%)	16.0	2.6	41.2 ²	33.8 ²	5.1

¹ not all degrees listed, ² physical activity data from Norway and 47% from Sweden are missing, in Norway calcium intake was low, and in Sweden and Norway vitamin D intake high, thus missing categories differ strongly between the quintiles; ³ Dietary intake from FFQ data, FAMS = monounsaturated fatty acids, FAPU = polyunsaturated fatty acids, FASAT = saturated fatty acids

Table 2: Dietary vitamin D and calcium intake in the respective countries of the EPIC cohort

	N, total cohort	Follow-Up time in person years	N, breast cancer cases	Vitamin D intake (µg/day) Mean ¹	Calcium intake (mg/day) Mean ¹
Denmark	28,694	215,799	822	4.67	1010
France	67,696	739,238	2844	2.68	907
Germany	27,864	227,045	455	3.63	959
Italy	30,434	256,918	673	2.08	818
Norway	35,226	210,297	470	4.15	808
Spain	24,809	241,082	316	5.00	981
Sweden	26,287	270,596	679	7.15	938
The Netherlands	26,432	228,577	564	4.31	1046
United Kingdom	52,543	441,127	937	3.74	968

¹ dietary intake from foods; data derived from 24-h diet recalls obtained in a subcohort (N = 36,994)

Table 3: Dietary vitamin D and calcium intake and risk of pre- and postmenopausal breast cancer

Vitamin D (µg/day)			All women						Premenopausal women			Postmenopausal women		
			Model 1 ¹			Model 2 ²			Model 2 ²			Model 2 ²		
			HR	95 % CI		HR	95 % CI		HR	95 % CI		HR	95 % CI	
Quintiles (range)	Median intake	Cases (N)												
Q1 (<1.85)	1.35	1563	1.00			1.00			1.00			1.00		
Q2 (1.85 - <2.69)	2.27	1640	0.95	0.88	1.02	0.94	0.87	1.01	0.98	0.85	1.13	0.90	0.82	0.99
Q3 (2.69 - <3.68)	3.15	1673	1.02	0.95	1.10	1.01	0.94	1.09	1.14	0.99	1.32	0.97	0.87	1.07
Q4 (3.68 - <5.46)	4.39	1531	1.00	0.93	1.09	0.99	0.91	1.08	1.04	0.88	1.23	0.99	0.89	1.11
Q5 (e 5.46)	7.35	1353	1.03	0.94	1.13	1.04	0.94	1.14	1.07	0.87	1.32	1.02	0.90	1.16
p for trend ³			0.92			0.92			0.78			0.21		
Calcium (mg/day)														
Quintiles (range)														
Q1 (<635)	516	1281	1.00			1.00			1.00			1.00		
Q2 (635 - <815)	730	1543	1.02	0.94	1.10	1.00	0.92	1.08	0.97	0.82	1.13	1.01	0.91	1.12
Q3 (815 - <992)	900	1575	0.96	0.88	1.04	0.93	0.86	1.01	0.92	0.77	1.09	0.94	0.84	1.05
Q4 (992 - <1,231)	1,097	1640	0.94	0.87	1.03	0.92	0.84	1.00	0.85	0.70	1.02	0.95	0.85	1.07
Q5 (e 1,231)	1,440	1721	0.93	0.85	1.02	0.91	0.83	1.01	0.98	0.80	1.19	0.90	0.79	1.02
p for trend ³			0.05			0.06			0.85			0.05		

¹ Cox proportional hazard model stratified by age and center adjusted for energy intake (kcal/day, quintiles)

² Cox proportional hazard model stratified by age and center, adjusted for no-fat, no-alcohol energy (kcal/day, quintiles), fat (g/day, quintiles, monounsaturated, polyunsaturated, and saturated fat), alcohol consumption (g/day, quintiles), weight (kg, cont.), height (cm, cont.), smoking status (never, former, current, unknown), education level (none or primary school, technical/professional school, secondary school, longer education incl. university degree, unknown), menopausal status (premenopausal, postmenopausal, perimenopausal), current use of contraceptives or hormones (yes, no, unknown), physical activity (occupational, recreational and household activity; inactive, moderately inactive, moderately active, active, missing), age at menarche (d12, 13-14, e15, missing)

³ Wald statistic of the continuous variable.

HR=hazard ratio, CI=confidence interval, 1 µg vitamin D = 40 IU vitamin D